



INOQ Rhodazo

Charge IFP eri T/11

Garden and Landscaping, especially for Ericads

Blends perfectly with all substrates
Increases absorption of water
Improves soil structure and aggregation
Increases humus content
May be used in organic farming in accordance with EC Commission Regulation 889/2008 Article 3 (4)

Ericomycorrhizal Fungi

(native strains, do not contain Genetically Modified Organisms (GMOs))

Hymenoscyphus ericae

Oidiodendron spp.

Mycorrhiza Effectiveness

(increased growth [%] in standardized test)

24 ± 6

Carrier material

Peat substrate

Specific weight [g/l]

250-400

pH

5,0

Application

Annual application near root system

Dosage

Blending into substrat

2 to 5 % (seedling production)
5 to 10 % (plant production)

Spreading into planting hole

20 ml / plant (up to 12 cm pot)
up to 100 ml / plant (up to 10 l pot)

Existing plant formations

According to size of plant up to
100 ml / plant in drilling holes

Surface treatment

Up to 100 ml / m²

Packaging

25 l PP-Bag, 1000 l Big Bag

Storage

6 months, under cool and dry conditions

Tolerance of fungicides

proven

Health and safety information

No special precaution necessary
Avoid breathing or ingestion
Absence of phytopathogens proven
(DNA multiscan®)
Material Safety Data Sheet available

For further technical information on application and products please contact:
INOQ GmbH, Geschäftsführung: Dr. Carolin Schneider, Solkau 2, 29465 Schnega
Tel. 0 58 42/98 16 72, Fax. 0 58 42/4 93, info@inoq.de · www.inoq.de